




Late Mississippian Gastropods of the Chainman Shale, West-Central Utah: Usgs Professional Paper 1368

By Mackenzie Gordon

BiblioGov. Paperback. Book Condition: New. This item is printed on demand. Paperback. 140 pages. Dimensions: 9.7in. x 7.4in. x 0.3in. The Chainman Shale of Mississippian (Osagean to late Chesterian) age, well exposed in the Confusion Range of western Utah, has yielded a profusion of fossils during investigations conducted by the U. S. Geological Survey in the past 30 years. Conspicuous among these fossils are gastropods, which range in age from latest Meramecian to late Chesterian. In west-central Utah, not far from the State boundary, the Chainman outcrop belt stretches from Granite Mountain south to the northern part of the Needle Range, a distance of69 miles (110 km). The Chainman thickens from north to south; the section at Granite Mountain is 1, 315 feet (401 m) thick and that at Jensen Wash in the Burbank Hills, 2, 203 feet (671 m). The rocks of the Chainman Shale record a general though irregular shallowing of the area from moderate depths of 330 feet (100 m) or so to quite shallow depths of perhaps locally little more than 3-6 feet (1-2 m). Most of the gastropods occur with ammonoids in a facies of shale or shale containing phosphatic limestone concretions, In this lutaceous facies,...



Reviews

An exceptional book and also the font utilized was intriguing to read. This is for all who statte there was not a worth reading. It is extremely difficult to leave it before concluding, once you begin to read the book.

-- Prof. Tyson Hilpert

This publication will never be effortless to begin on studying but extremely entertaining to learn. It is probably the most incredible publication i have go through. I realized this ebook from my i and dad suggested this publication to learn.

-- Austin O'Connell